


# Quantization (Jan 22, 2004)

---

- Optimal Quantizer
- Uniform Quantizer

# Image Quantization



# Decision/Reconstruction

## Levels

$$u \in [t_k, t_{k+1}] \longrightarrow r_k$$

$\{t_k : k=1,2,\dots,L+1\}$        $\longleftarrow$       Transition or decision levels

$$r_k \longrightarrow k^{\text{th}} \text{ reconstruction level}$$

**Example:** Uniform quantizer  $u \in [0,10.0]$

We want  $u' \in \{0,1,\dots,255\}$

$t_1 = 0; t_{257} = 10.0$ ; uniformly spaced,  $t_k = (k-1).10/256$

$$k = 1,2,\dots,257)$$

# Example: quantization

---

$$r_k = t_k + \frac{1}{2} \left( \frac{10}{256} \right) = t_k + \frac{5}{256}$$

Quantization interval

$$q_k = t_k - t_{k-1} = r_k - r_{k-1}$$

= Constant  $\Rightarrow$  Uniform quantizer

# MMSE Quantizer

---

**Minimise the mean squared error,  $MSE =$   
Expected value of  $(u-u')^2$  given the number of  
quantization levels  $L$ .**

Assume that the density function  $p_u(u)$  is known  
(or can be approximated by a normalised  
histogram).

Note that for images,  $u$ ==image intensity.  $p_u(u)$  is  
the image intensity ditribution.

# Optimum MSE quantizer

---

$$E(u) = \text{Expected value of } u = \int_{-\infty}^{\infty} u p_u(u) du$$

$$MSE, \epsilon = E((u - u')^2) = \int_{-\infty}^{\infty} (u - u')^2 p_u(u) du = \int_{t_1}^{t_{L+1}} (u - u')^2 p_u(u) du$$

Since  $u' = r_k$  if  $u \in [t_k, t_{k+1}]$ , we can rewrite this as

$$\epsilon = \sum_{i=1}^L \int_{t_i}^{t_{i+1}} (u - r_i)^2 p_u(u) du$$

Conditions for minimisation of  $\epsilon$  are:  $\frac{\partial \epsilon}{\partial r_k} = 0$ ;  $\frac{\partial \epsilon}{\partial t_k} = 0$

# MMSE (contd.)

---

$$\frac{\partial \mathcal{E}}{\partial t_k} = (t_k - r_{k-1})^2 p_u(t_k) - (t_k - r_k)^2 p_u(t_k) = 0$$

$$\text{Now } t_k \leq r_k < t_{k+1} \Rightarrow t_k - r_{k-1} = r_k - t_k \Rightarrow \left[ t_k = \frac{(r_k + r_{k-1})}{2} \right] \dots \text{(A)}$$

$$\frac{\partial \mathcal{E}}{\partial r_k} = \int_{t_k}^{t_{k+1}} 2(u - r_k) (-1) p_u(u) du = 0$$

$$\begin{aligned} & \int u p_u(u) du \\ \Rightarrow r_k &= \frac{\int_{t_k}^{t_k} u p_u(u) du}{\int_{t_k}^{t_{k+1}} p_u(u) du} = \text{E}(u \mid u \in [t_k, t_{k+1}]) \end{aligned} \quad \dots \text{(B)}$$

# Optimum transition/reconst.

---

- (1) Optimal transition levels lie halfway between the optimum reconstruction levels.
- (2) Optimum reconstruction levels lie at the center of mass of the probability density in between the transition levels.
- (3) A and B are simultaneous non-linear equations (in general)  
→ Closed form solutions normally don't exist → use numerical techniques

# Uniform optimal quantizer

---

Consider  $p_u(u) = \begin{cases} \frac{1}{t_{L+1} - t_1} & t_1 \leq u \leq t_{L+1} \\ 0 & \text{Otherwise} \end{cases}$

Then  $r_k = \frac{\int_{t_k}^{t_{k+1}} u p_u(u) du}{\int_{t_k}^{t_{k+1}} p_u(u) du} = \frac{\frac{1}{t_{L+1} - t_1} \int_{t_k}^{t_{k+1}} u du}{\frac{1}{t_{L+1} - t_1} \int_{t_k}^{t_{k+1}} du} = \frac{\left(\frac{u^2}{2}\right)_{t_k}^{t_{k+1}}}{(u)_{t_k}^{t_{k+1}}} = \frac{(t_{k+1}^2 - t_k^2)}{2(t_{k+1} - t_k)}$

$$r_k = \frac{1}{2}(t_{k+1} + t_k); \quad t_k = \frac{r_k + r_{k-1}}{2} = \frac{1}{2} \cdot \frac{1}{2} [t_{k+1} + t_k + t_k + t_{k-1}] = \frac{t_{k+1} + t_{k-1}}{2}$$

# Uniform Quantizer

---

$$t_k - t_{k-1} = t_{k+1} - t_k = \text{Constant} \quad q = \frac{t_{L+1} - t_1}{L}$$

$$t_k = t_{k-1} + q \quad ; \quad r_k = t_k + \frac{q}{2}$$

Quantization error  $e = (u - u')$  is uniformly distributed over the interval

$$\left( -\frac{q}{2}, \frac{q}{2} \right)$$

$$\text{Mean squared error } E((u - u')^2) = \frac{1}{q} \int_{-\frac{q}{2}}^{\frac{q}{2}} u^2 = \frac{q^2}{12}$$